

Operator fidelity susceptibility, decoherence and quantum criticality

Xiao-Ming Lu,¹ Zhe Sun,^{1,2} Xiaoguang Wang,^{1,*} and Paolo Zanardi^{3,4}

¹*Zhejiang Institute of Modern Physics, Department of Physics, Zhejiang University, Hangzhou 310027, China*

²*Department of Physics, Hangzhou Normal University, Hangzhou 310036, China*

³*Department of Physics and Astronomy, University of Southern California, Los Angeles, CA 90089-0484 (USA)*

⁴*Institute for Scientific Interchange, Viale Settimio Severo 65, I-10133 Torino, Italy*

The extension of the notion of quantum fidelity from the state-space level to the operator one can be used to study environment-induced decoherence. State-dependent operator fidelity susceptibility (OFS), the leading order term for slightly different operator parameters, is shown to have a nontrivial behavior when the environment is at critical points. Two different contributions to OFS are identified which have distinct physical origins and temporal dependence. Exact results for the finite-temperature decoherence caused by a bath described by the Ising model in transverse field are obtained.

I. INTRODUCTION

Generically the interaction between a quantum system and its environment results in decoherence, and may lead the system to experience the so-called quantum-classical transition [1]. For this reason the decoherence process is regarded as the main obstacle for the implementation of quantum information processing [2]. Generally speaking the properties of the environment may strongly affect its decohering capabilities [3, 4, 5, 7, 8]. This implies that a quantum system can be regarded as a probe to extract useful information about the coupled environment, e.g., the quantum phase transitions (QPTs). Very recently a nuclear magnetic resonance (NMR) experiment of this kind has been performed [9]. There a single qubit was used as a probe to detect the (precursors of the) quantum critical point of the coupled environment.

The relationship between the decoherence of a central spin and the QPTs of the coupled environment can be established through the notion of Loschmidt echo (LE) [5]. It is well known that LE can be exploited to measure the stability of a quantum system to perturbations [3, 10, 11]. In Ref. [5], a central spin coupled with an Ising spin chain in a transverse field was considered; the authors found that the decay of LE is enhanced by quantum criticality. The connection between LE and quantum criticality has been further clarified in Ref. [6]. There the authors showed that the LE enhancement holds true just for short decay times (gaussian regime). A genuine signature of criticality, on the other hand, can be recovered by studying the asymptotic (large time) behaviour of LE as function of the environment coupling parameter [6]. More recently, the averaged LE over all states on Hilbert space with a Haar measure, called operator fidelity, was proposed to study quantum criticality [12]. This proposal is a direct extension to the operator level of state-space quantum fidelity. This latter notion recently attracted a lot of attention as a new tool to ana-

lyze quantum criticality, both a zero-temperature and at finite temperature [13, 14, 15, 16, 17, 18].

Whereas in [5] the initial state of the spin chain was assumed to be the ground state, in this paper we aim at considering a more general situation: the initial state of the environment is a mixture of the eigenstate of its Hamiltonian. The Gibbs thermal state is special instance of this setup. With this assumption the study of decoherence naturally leads to a state-dependent generalizations of the operator fidelity suggested in [12].

In this paper we will discuss the operator extensions of fidelity on general grounds. In order to do that one has simply to notice that finite-time quantum evolutions correspond to unitary operators that themselves belong to an Hilbert space i.e., the Hilbert-Schmidt one. It follows that some of the results obtained in the fidelity approach for quantum states lifts immediately to the operator level we are now interested in. We will show that the corresponding operator fidelity susceptibility (OFS) contains two different contributions. These two terms arise from variation upon parameter change of energy levels and eigenstates respectively and have very different temporal behaviour. We will then get a general expression for OFS for models with a factorization structure typical of quasi-free models. Finally, we will exploit OFS for studying an environment described by an Ising chain with a transverse field that can be driven to quantum criticality.

This paper is organized as follows: In Sec. II, we give a general description of the models and the relations between decoherence of the central system and the operator fidelity of two time evolutions of the environment. In Sec. III, we give generalities about operator metric and fidelities and analyze the general OFS. In Sec. IV A, we consider the models with a factorization structure, and give a general expression for the OFS, and then study the Ising model with a transverse field, get the exact solution for the OFS. Sec. V contains the conclusions and outlook.

*Electronic address: xgwang@zimp.zju.edu.cn

II. DECOHERENCE AND OPERATOR FIDELITY

As mentioned in the introduction, an important physical motivation for the operator fidelities we are going to analyze in this paper is provided by the decoherence process. To see this clearly, let us consider a purely dephasing coupling of a system with its environment. The interaction Hamiltonian has the form

$$H_I = \sum_n |n\rangle\langle n| \otimes B_n. \quad (1)$$

The entire Hamiltonian is $H = H_0 + H_I$, where H_0 consists of system Hamiltonian $H_S = \sum_n E_n |n\rangle\langle n|$ and bath Hamiltonian H_B . The H_S eigenstates $\{|n\rangle\}$, play here the role of preferred pointer states [1].

With the initial state $\rho(0) = |\psi(0)\rangle\langle\psi(0)| \otimes \rho_B$, the reduced density matrix of the system at time t can be written as

$$\rho_S(t) = \sum_{n,m} c_n c_m^* e^{-i(E_n - E_m)t} |n\rangle\langle m| \text{Tr}_B[\rho_B V_m^\dagger(t) V_n(t)], \quad (2)$$

where $c_n = \langle n|\psi(0)\rangle$, $V_n(t) = \exp[-it(H_B + B_n)]$

Obviously, in the generic case ρ_S evolves from initial pure states to mixed states. The decay of the off-diagonal elements of ρ_S means a reduction from a pure state to a classical mixture of the preferred pointer states — quantum-classical transitions. The temporal behavior of the off-diagonal element is decided by two factors, one is $c_n c_m^*$, just relating to initial state, and the other $\text{Tr}_B[\rho_B V_m^\dagger(t) V_n(t)]$ is unrelated with the initial state of the system, reflecting the dephasing effect induced by environment. The latter can be considered as a fidelity for two operators V_m and V_n . which is defined by

$$F_\rho(V_m, V_n) = |\langle V_m, V_n \rangle_\rho| \quad (3)$$

where the inner product $\langle V_m, V_n \rangle_\rho = \text{Tr}[\rho V_m^\dagger V_n]$. This fidelity can be obtained from the density matrix of the central system and encodes information about the bath state. Notice that one might choose $B_n = \lambda_n B$; in this case different V_n simply correspond to different values of the coupling strength λ in front of the "perturbation" B . This is the scenario we mostly have in mind in this paper (see Sec. IV).

Let us now specialize to a 2-level system coupled with a bath. If interaction is weak enough, the two effective Hamiltonians are slightly different. For an instance in Sec. IV, we will consider the important case where the initial state of the bath is a thermal equilibrium state $\rho_B = \exp(-\beta H)/Z(\beta)$ [$Z(\beta) = \text{Tr}[\exp(-\beta H)]$ is the partition function, $T = \beta^{-1}$ is the temperature of the bath]. In such a case $|F_{\rho_B}(V_1, V_2)|^2$ will become the Loschmidt echo for $\beta^{-1} = 0$ (for non degenerate ground state) and it coincides with the operator fidelity defined in [12] for $\beta = 0$.

III. METRICS OVER MANIFOLDS OF UNITARIES

In this section we discuss operator fidelity from a rather general mathematical point view. Let $\mathcal{H}$ be a quantum state-space and a $\{U_\lambda\} \subset \mathcal{U}(\mathcal{H})$ a smooth family of unitaries over $\mathcal{H}$ parametrized by elements of a manifold $\mathcal{M}$. Given a state $\rho \in \mathcal{S}(\mathcal{H}) = \{\rho \in \mathcal{L}(\mathcal{H}) | \rho \geq 0, \text{tr} \rho = 1\}$ one can define the following hermitean product over $\mathcal{L}(\mathcal{H})$

$$\langle X, Y \rangle_\rho := \text{Tr}(\rho X^\dagger Y) \quad (4)$$

If $\rho > 0$ then (4) is non-degenerate and $\|X\|_\rho := \sqrt{\langle X, X \rangle_\rho}$ defines a norm over $\mathcal{L}(\mathcal{H})$. In general, if $\ker \rho \neq \{0\}$, $\|\bullet\|_\rho$ is just a semi-norm (if the range of X is included in $(\text{supp} \rho)^\perp$ then $\|X\|_\rho = 0$) Notice that i) $\|X\|_\rho \leq \|X\|$ and ii) unitaries U are normalized i.e., $\langle U, U \rangle_\rho = 1$.

Definition The ρ -fidelity of the operators X and Y is given by

$$\mathcal{F}_\rho(X, Y) := |\langle X, Y \rangle_\rho| \quad (5)$$

It is immediate yet important to realize that for non full-rank states ρ , having two unitaries with ρ -fidelity one does not imply their equality (up to a phase). Indeed, from the Cauchy-Schwarz inequality one has $\mathcal{F}_\rho(X, Y) \leq \|X\|_\rho \|Y\|_\rho$, in particular for unitaries U and W , $\mathcal{F}_\rho(U, W) \leq 1$. One has that $\mathcal{F}_\rho(U, W) = 1 \Leftrightarrow U|_{\text{supp} \rho} = W|_{\text{supp} \rho}$ where $\text{supp} \rho := (\ker \rho)^\perp$.

In order to unveil the operational meaning of the above definition it is useful to recall the following fact. If $|\Psi_\rho\rangle \in \mathcal{H} \otimes \mathcal{H}$ is a purification of $\rho = \sum_i p_i |i\rangle\langle i|$ i.e., $|\Psi_\rho\rangle = \sum_i \sqrt{p_i} |i\rangle \otimes |i\rangle$ then $\langle X, Y \rangle_\rho = \langle \psi_\rho | (X^\dagger \otimes \mathbb{1})(Y \otimes \mathbb{1}) | \Psi_\rho \rangle$. The operator scalar product (4) can be seen as a scalar product of suitable quantum states of a bigger system. This simple remark shows that operator fidelity (5) quantifies the degree of statistical distinguishability between the two states

$$|\Psi(A)\rangle := (A \otimes \mathbb{1}) |\Psi_\rho\rangle \quad (A = X, Y).$$

When U and W denote unitary transformations the quantity (5) has an interpretation as visibility strength in properly designed interferometric experiments [25]. Yet another kind of operational relevance of the operator fidelities in the context of decoherence will be discussed in II.

Now, following the differential-geometric spirit of [20], we are going to consider the operator fidelity (5) between infinitesimally different unitaries. The leading term in the expansion of (5) will define a quadratic form over the tangent space of the manifold $\mathcal{U}(\mathcal{H})$. For full rank ρ that quadratic form is a *metric*. The following proposition shows that and its proof it is just a direct calculation analogous to the one performed at the state-space level [20].

Proposition Let $\{U_\lambda\} \subset \mathcal{U}(\mathcal{H})$ be a smooth family of unitaries over $\mathcal{H}$ parametrized by elements λ of a manifold $\mathcal{M}$. One finds

$$\mathcal{F}_\rho(U_\lambda, U_{\lambda+\delta\lambda}) = 1 - \frac{\delta\lambda^2}{2} \chi_\rho(\lambda) \quad (6)$$

where, if $U' = \partial U / \partial \lambda$ one has

$$\chi_\rho(\lambda) := \langle U', U' \rangle_\rho - |\langle U', U \rangle_\rho|^2 \quad (7)$$

The quantity above will be referred to as operator fidelity susceptibility (OFS). In the following we will be discussing three different situations fitting in the framework we have set. The first example (**Ex. 0**) will show in which precise sense the operator approach here under investigation includes the ground-state fidelity one. In **Ex. 1** we notice that when the family of unitaries is a one-parameter group generated by an Hamiltonian H then the metric is nothing but the variance of H . Finally, in **Ex. 2** the most important case of a fixed-time family of evolutions generated by a parametric family of Hamiltonians is analyzed.

Example 0 Let $\{H_\lambda\}$ a family of *isodegenerate* quantum Hamiltonians with spectral resolutions

$$H(\lambda) = \sum_n E_n(\lambda) |\psi_n(\lambda)\rangle \langle \psi_n(\lambda)|.$$

The adiabatic intertwiners are unitaries such that $O(\lambda, \lambda_0) |\psi_n(\lambda_0)\rangle = |\psi_n(\lambda)\rangle \forall n$. Let $\rho = |\psi_0(\lambda_0)\rangle \langle \psi_0(\lambda_0)|$ ($|\psi_0(\lambda_0)\rangle$ = Ground State of $H(\lambda_0)$) and $U(\lambda) = O(\lambda, \lambda_0)$ then

$$\langle U(\lambda), U(\lambda') \rangle_\rho = \langle \psi_0(\lambda_0) | O^\dagger(\lambda, \lambda_0) O(\lambda', \lambda_0) | \psi_0(\lambda_0) \rangle = \langle \psi_0(\lambda), \psi_0(\lambda') \rangle. \quad (8)$$

This shows that the ground state fidelity is a particular instance of the setting we are discussing.

Example 1 Let $\{U_t = e^{-itH}\}_t$ and $\rho = |\phi\rangle \langle \phi|$. Then $U' = -iHU(t)$ whence $\chi_\rho(t) = \langle \phi | H^2 | \phi \rangle - |\langle \phi | H | \phi \rangle|^2$

Example 2 $\{U_\lambda = e^{-itH_\lambda}\}_\lambda$ and $\rho = \mathbb{1}/d$ ($d := \dim \mathcal{H}$). From time-dependent perturbation theory one obtains

$$U' = -iU_\lambda \int_0^t d\tau e^{i\tau H_\lambda} H' e^{-i\tau H_\lambda} \quad (9)$$

Introducing the superoperator $L_H := [H, \bullet]$ over $\mathcal{L}(\mathcal{H})$ one sees that, if $|n\rangle$'s and the E_n 's denote the eigenvectors and eigenvalues of $H(\lambda)$, $E_n - E_m$ and $\hat{\Psi}_{n,m} := \sqrt{d}|n\rangle \langle m|$ are the eigenvalues and (normalized) eigenoperators of L_H , respectively. One has $\langle U', U' \rangle = \langle \int_0^t d\tau e^{i\tau L_H} (H'), \int_0^t d\sigma e^{i\sigma L_H} (H') \rangle$. If $P = \sum_n |\hat{\Psi}_{n,n}\rangle \langle \hat{\Psi}_{n,n}|$ is the projection over the kernel of L_H and $Q = 1 - P$ one finds $e^{i\tau L_H} = Q e^{i\tau L_H} Q + P$, therefore the above expression becomes

$$\begin{aligned} \langle U', U' \rangle &= \langle Q \frac{e^{i\tau L_H} - 1}{L_H} Q (H'), Q \frac{e^{i\tau L_H} - 1}{L_H} Q (H') \rangle + \\ &+ t^2 \langle P (H'), P (H') \rangle = \|F_t Q (H')\|^2 + t^2 \|P (H')\|^2 \end{aligned} \quad (10)$$

(notice that L_H is invertible, by definition, on the range of Q) where $F_t(x) := \sin(xt/2)/(x/2)$.

The first term in the last line of (10) can be rewritten as $\langle H' | Q F_t^2(L_H) Q | H' \rangle$. Similarly the term $\langle U, U' \rangle$

in the metric can be written as $\langle 1, \int_0^t d\tau e^{i\tau L_H} (H') \rangle = \int_0^t d\tau \langle e^{-\tau L_H} (1), H' \rangle = t \langle 1, H' \rangle = d^{-1} \text{Tr} H'$. Putting all together and expanding over the basis $|\hat{\Psi}_{n,n}\rangle$ of L_H -eigenoperators one finds the explicit result

$$\chi_\rho(\lambda) = \frac{1}{d} \sum_{n \neq m} \left| \frac{\langle n | H' | m \rangle}{E_n - E_m} \right|^2 F_t^2(E_n - E_m) \quad (11)$$

$$+ t^2 \left(\frac{1}{d} \sum_n |\langle n | H' | n \rangle|^2 - \frac{1}{d^2} |\text{Tr} H'|^2 \right) \quad (12)$$

The key property here is that

$$\lim_{t \rightarrow \infty} t^{-1} F_t^2(x) = 2\pi \delta(x).$$

This asymptotic delta function is responsible for the large contribution to (12) given by small $E_n - E_m$. This shows that *all* (quasi) level crossings in the spectrum of H might lead to a analyticity breakdown in χ_ρ . This has to be contrasted with the GS fidelity studied e.g., in [13] where just (quasi) level crossings in the GS play a role. In this sense it is clear why the OFS (12) is a more powerful tool than the corresponding state-space analogue.

An important generalization of (12) can be obtained by considering ρ *commuting* with H . In this case one obtains an expression analogous to (12) with the diagonal elements $\rho_{n,n} = \langle n | \rho | n \rangle$ suitably inserted (see Sec. III A).

A. Splitting χ_ρ

Now we derive an alternative form for the OFS (7) for the case of *Ex. 2* discussed above. This form will make even more manifest the different contributions to operator susceptibility arising from eigenvalue and eigenvector variation with the control parameter λ . A unitary operator can be written in diagonal form

$$U(\lambda) = \sum_i u_i(\lambda) P_i(\lambda). \quad (13)$$

where the u_i 's are the eigenvalues of U satisfying $|u_i| = 1$ and $P_i = |\psi_i\rangle \langle \psi_i|$ is an one-dimensional projective operator. Both eigenvalues and eigenstates are parameter dependent. What we would like to do first is to separate the contributions of these two kinds of parameter dependence to the susceptibility $\chi_\rho(\lambda)$. In Ref. [16], the authors succeed in telling apart the classical and quantum contribution to the Bures metric with consideration of these two kinds of dependence.

The differential of this unitary operator can be divided into two parts

$$\begin{aligned} \partial_\lambda U(\lambda) &= D_1(\lambda) + D_2(\lambda), \\ D_1(\lambda) &= \sum_i (\partial_\lambda u_i) P_i, \quad D_2(\lambda) = \sum_i u_i (\partial_\lambda P_i), \end{aligned} \quad (14)$$

where $\partial_\lambda P_i = |\psi_i\rangle\langle\partial_\lambda\psi_i| + |\partial_\lambda\psi_i\rangle\langle\psi_i|$. Next, we assume that the density matrix ρ can be simultaneously diagonalized with U , so it can be written as the form

$$\rho = \sum_i \rho_{ii} P_i. \quad (15)$$

The assumption $[H, \rho] = 0$ is motivated by the important example where both time evolution unitary operators and density operators considered are generated by the same Hamiltonian i.e., ρ is the Gibbs thermal state associated with H .

Since

$$\langle P_j \partial_\lambda P_i \rangle_\rho = \delta_{ij} \rho_{jj} (\langle \psi_j | \partial_\lambda \psi_i \rangle + \langle \partial_\lambda \psi_i | \psi_j \rangle) = 0, \quad (16)$$

after substituting Eq. (14) into Eq. (7), we obtain

$$\chi_\rho(\lambda) = \chi_\rho^{(1)}(\lambda) + \chi_\rho^{(2)}(\lambda), \quad (17)$$

$$\begin{aligned} \chi_\rho^{(1)}(\lambda) &= \langle D_1^\dagger D_1 \rangle_\rho - |\langle U^\dagger D_1 \rangle_\rho|^2, \\ \chi_\rho^{(2)}(\lambda) &= \langle D_2^\dagger D_2 \rangle_\rho. \end{aligned} \quad (18)$$

Here we separate the contributions of D_1 and D_2 . To make them explicit, we consider the time evolution operator $U(\lambda, t) = e^{-itH(\lambda)}$, and assume that the eigenstates changing with λ can be connected through a smooth unitary transformation $\mathcal{S}(\lambda)$ which is time-independent and satisfies $\mathcal{S}(\lambda)H(\lambda)\mathcal{S}(\lambda)^\dagger = H_d(\lambda)$, where H_d is a diagonal Hamiltonian in the fixed λ -independent basis. Therefore, the unitary operators considered can be written

$$U(\lambda, t) = \mathcal{S}^\dagger(\lambda) U_d(\lambda, t) \mathcal{S}(\lambda), \quad (19)$$

with $U_d = \exp(-itH_d)$. After differentiating with respect to λ , we get

$$\begin{aligned} D_1(\lambda, t) &= -it\mathcal{S}^\dagger(\lambda)[\partial_\lambda H_d(\lambda)]\mathcal{S}(\lambda)U(\lambda, t), \\ D_2(\lambda, t) &= [\mathcal{A}(\lambda), U(\lambda, t)], \end{aligned} \quad (20)$$

where

$$\mathcal{A}(\lambda) = (\partial_\lambda \mathcal{S}(\lambda))^\dagger \mathcal{S}(\lambda). \quad (21)$$

Substituting Eq. (21) into Eq. (18) leads to

$$\chi_\rho^{(1)}(\lambda, t) = t^2 [\langle (\partial_\lambda H_d)^2 \rangle_\rho - \langle \partial_\lambda H_d \rangle_\rho^2]. \quad (22)$$

This can be considered as the fluctuation of the quantity $\partial_\lambda H_d(\lambda)$ under the state ρ , with an extra factor t^2 .

The second term of Eq. (17) will be

$$\chi_\rho^{(2)}(\lambda, t) = \langle [U^\dagger(\lambda, t), \mathcal{A}^\dagger(\lambda)] [\mathcal{A}(\lambda), U(\lambda, t)] \rangle_\rho. \quad (23)$$

Since U and ρ can be diagonalized simultaneously, the above expression can be further rewritten as $\chi_\rho^{(2)}(\lambda, t) = 2 \sum_{mn} \rho_{nn} |\mathcal{A}_{mn}|^2 (1 - \cos[(E_n - E_m)t])$. This form explicitly shows that the time-dependent terms in $\chi_\rho^{(2)}(\lambda, t)$ are circular functions.

Now we would like to make a few general comments about Eqs. (22) and (23):

- i) They correspond one to one to the two terms in Eq. (12) [with suitably inserted $\rho_{n,n}$]. In particular this remark shows that $\mathcal{A}_{n,m} = (E_n - E_m)^{-1} \langle n | H' | m \rangle$. Notice that $\mathcal{A}$ is nothing but the infinitesimal generator of the adiabatic intertwiner mentioned in Sec III.
- ii) $\chi_\rho^{(i)}$ ($i = 1, 2$) separate apart the contributions of the eigenvalue variations from those of the eigenstates. In Ref. [16], a similar distinction was made for the Bures metric on thermal state manifolds (the corresponding terms there were named the classical and the quantum, respectively).
- iii) Eqs (22) and (23) have quite different forms of time dependence. $\chi_\rho^{(1)}$ gives t^2 contributions explicitly, while the second term consists of circular functions with finite period of criticality. When t is large, if $\chi_\rho^{(1)}$ is not vanishing, it dominates the OFS and therefore the decay behavior of the operator fidelity.
- iv) If ρ is pure [implying, since we have assumed the form $\rho = \sum_i \rho_{ii} P_i$, that the initial state is an eigenstate of the Hamiltonian] $\chi_\rho^{(1)}$ vanishes. Moreover if $\mathcal{A}(\lambda)$ commutes with $U(\lambda, t)$, and therefore the eigenvectors of $U(\lambda)$ are λ -independent, it is obvious from (23) that $\chi_\rho^{(2)}$ vanishes.

IV. THE TRANSVERSE FIELD ISING MODEL

In this section, we apply the general formalism developed so far to the concrete case of the transverse field Ising mode. Before doing that it is useful to discuss a bit generally models having a factorization structure.

A. Factorizable models

In order to obtain the exact solution of $\chi_\rho(\lambda, t)$, we analyze the cases where the unitary operator $U(\lambda, t)$ and thermal state $\rho(\lambda, \beta)$ have the same factorization structure. This assumption hold true, for example, in cases where they are both generated by the same Hamiltonian $H(\lambda)$. These two quantities can be then written in the composite space

$$\begin{aligned} U(\lambda) &= \bigotimes_{k=0}^M U_k(\lambda), \\ \rho(\lambda) &= \bigotimes_{k=0}^M \rho_k(\lambda), \end{aligned} \quad (24)$$

where U_k is still an unitary operator and ρ_k is still a density operator, corresponding to the k -th subspace. Note that

$$\partial_\lambda U(\lambda) = \sum_{l=0}^M \left[\bigotimes_{k=0}^{l-1} U_k(\lambda) \right] \otimes \partial_\lambda U_l(\lambda) \otimes \left[\bigotimes_{k=l+1}^M U_k(\lambda) \right]. \quad (25)$$

Since $\text{Tr}(A \otimes B) = \text{Tr}(A)\text{Tr}(B)$ and $\text{Tr}(\rho_l) = 1$, the first and second terms of Eq. (7) can be expressed respectively as

$$\begin{aligned} \text{Tr}[\rho(\partial_\lambda U)^\dagger(\partial_\lambda U)] &= \sum_l \text{Tr}_l[\rho_l(\partial_\lambda U_l)^\dagger(\partial_\lambda U_l)] + \sum_{l \neq l'} A_{ll'} \\ |\text{Tr}[\rho U^\dagger(\partial_\lambda U)]|^2 &= \sum_l |\text{Tr}_l[\rho_l U_l^\dagger(\partial_\lambda U_l)]|^2 + \sum_{l \neq l'} A_{ll'} \end{aligned} \quad (26)$$

where $A_{ll'} = \text{Tr}_l[\rho_l U_l^\dagger(\partial_\lambda U_l)]\text{Tr}_{l'}[\rho_{l'} U_{l'}^\dagger(\partial_\lambda U_{l'})]$, and the subscript l means the l -th subspace.

Thus, the OFS χ_ρ for the factorized unitary operator is

$$\chi_\rho(\lambda) = \sum_l \chi_{\rho,l}(\lambda) \quad (27)$$

$$\begin{aligned} \chi_{\rho,l}(\lambda) &= \text{Tr}_l[\rho_l(\partial_\lambda U_l)^\dagger(\partial_\lambda U_l)] \\ &\quad - |\text{Tr}_l[\rho_l U_l^\dagger(\partial_\lambda U_l)]|^2 \end{aligned} \quad (28)$$

B. Ising Hamiltonian

We now consider transverse field Ising model. The Hamiltonian is given by

$$H(\lambda) = -J \sum_{l=-M}^M (\sigma_l^x \sigma_{l+1}^x + \lambda \sigma_l^z), \quad (29)$$

where J is an exchange constant hereafter assumed to be the unity, and λ is the transverse field strength. This model can be calculated analytically by using the Jordan-Wigner transformation [22]

$$\begin{aligned} \sigma_i^x &= \prod_{j < i} (1 - 2c_j^\dagger c_j) (c_i + c_i^\dagger), \\ \sigma_i^z &= 1 - 2c_i^\dagger c_i, \end{aligned} \quad (30)$$

which maps the spins to fermions. After the Fourier transformations, the Hamiltonian in the momentum space is

$$\begin{aligned} H(\lambda) &= - \sum_{k=-M}^M [\cos(\frac{2\pi k}{N}) - \lambda] (d_k^\dagger d_k + d_{-k}^\dagger d_{-k} - 1) \\ &\quad + i \sin(\frac{2\pi k}{N}) (d_k^\dagger d_{-k}^\dagger - d_{-k} d_k), \end{aligned} \quad (31)$$

where $N = 2M + 1$. This Hamiltonian can be exactly solved by a Bogoliubov transformation [23]. However, introducing a set of pseudo-spin operators is more convenient here. Since $n_k - n_{-k}$ ($n_k = d_k^\dagger d_k$) commutes with every term of the Hamiltonian (31), the pseudo Pauli operators can be defined by [24]

$$\begin{aligned} \varsigma_{kx} &= d_k^\dagger d_{-k}^\dagger + d_{-k} d_k, \\ \varsigma_{ky} &= -i(d_k^\dagger d_{-k}^\dagger - d_{-k} d_k), \\ \varsigma_{kz} &= d_k^\dagger d_k + d_{-k}^\dagger d_{-k} - 1. \end{aligned} \quad (32)$$

These give the Pauli matrix in the $n_k - n_{-k} = 0$ subspace, and become zero matrix in the $n_k - n_{-k} = \pm 1$ subspaces. Therefore ς_{0x} , ς_{0y} , ς_{0z} are just the standard Pauli matrices σ_{0x} , σ_{0y} , and σ_{0z} .

With these operators, the Hamiltonian can be written as

$$\begin{aligned} H(\lambda) &= \sum_{k=1}^M \mathcal{S}_k^\dagger(\lambda) H_{k,d}(\lambda) \mathcal{S}_k(\lambda) + (\lambda - 1) \varsigma_{0z}, \\ H_{k,d} &= \Omega_k \varsigma_{kz}, \\ \mathcal{S}_k(\lambda) &= \exp(-i \frac{\theta_k}{2} \varsigma_{kx}), \end{aligned} \quad (33)$$

where

$$\begin{aligned} \Omega_k &= -2\sqrt{[\lambda - \cos(2\pi k/N)]^2 + \sin^2(2\pi k/N)}, \\ \theta_k &= \arcsin[\frac{2\sin(2\pi k/N)}{\Omega_k}]. \end{aligned} \quad (34)$$

We consider the fidelity of two time evolution operators $U(\lambda) = \exp(-itH(\lambda))$ and $U(\lambda + \delta\lambda)$ with thermal state $\rho = \exp(-\beta H(\lambda))/Z(\beta, \lambda)$, where $Z(\beta, \lambda) = \text{Tr}[\exp(-\beta H(\lambda))]$ is the partition function, and $\beta = 1/T$ is the inverse temperature. The operators U and ρ can be written in the factorized form

$$\begin{aligned} U &= e^{-it(\lambda-1)\sigma_{0z}} \otimes [\bigotimes_{k=1}^M U_k], \\ \rho &= \frac{e^{\beta(\lambda-1)\sigma_{0z}}}{Z_0} \otimes [\bigotimes_{k=1}^M \frac{\rho_k}{Z_k}], \end{aligned} \quad (35)$$

where

$$\begin{aligned} U_k &= \mathcal{S}_k^\dagger(\lambda) e^{-it\Omega_k \varsigma_{kz}} \mathcal{S}_k(\lambda) = e^{-it\Omega_k \varsigma_{kn}}, \\ \rho_k &= \mathcal{S}_k^\dagger(\lambda) e^{-\beta\Omega_k \varsigma_{kz}} \mathcal{S}_k(\lambda) = e^{-\beta\Omega_k \varsigma_{kn}}, \\ \varsigma_{kn} &= \varsigma_{ky} \sin \theta_k + \varsigma_{kz} \cos \theta_k, \end{aligned} \quad (36)$$

and the partition functions are

$$Z_0 = 2 \cosh(\beta(\lambda - 1)), \quad Z_k = 2[1 + \cosh(\beta\Omega_k)]. \quad (37)$$

For $k > 0$, after substituting $H_{k,d} = \Omega_k \varsigma_{kz}$ into Eq. (22), we have

$$\chi_{\rho,k}^{(1)}(\lambda, t) = 4t^2 \frac{\cos^2 \theta_k}{\cosh(\beta\Omega_k) + 1}. \quad (38)$$

To calculate $\chi_{\rho,k}^{(2)}$, we should first calculate the related quantities $\mathcal{A}_k$ defined by Eq. (21) and the commutator

$$\begin{aligned} \mathcal{A}_k &= i \frac{\theta'_k}{2} \varsigma_{kx}, \\ [\mathcal{A}_k, U_k] &= i \theta'_k \sin(t\Omega_k) (\varsigma_{kz} \sin \theta_k - \varsigma_{ky} \cos \theta_k), \end{aligned} \quad (39)$$

where θ'_k denotes $\partial_\lambda \theta_k$ for convenience. Substituting them into Eq. (23), we obtain

$$\chi_{\rho,k}^{(2)} = 4 \frac{\cosh(\beta\Omega_k)}{\cosh(\beta\Omega_k) + 1} \frac{\sin^2 \theta_k \sin^2(t\Omega_k)}{\Omega_k^2}. \quad (40)$$

For $k = 0$ subspace, the contribution is

$$\chi_{\rho,0}^{(1)} = t^2[1 - \tanh^2(\beta(\lambda - 1))], \quad \chi_{\rho,0}^{(2)} = 0. \quad (41)$$

Thus, we get the OFS

$$\begin{aligned} \chi_{\rho}^{(1)} &= t^2[1 - \tanh^2(\beta(\lambda - 1))] + 4 \sum_{k=1}^M t^2 \frac{\cos^2 \theta_k}{\cosh(\beta\Omega_k) + 1}, \\ \chi_{\rho}^{(2)} &= 4 \sum_{k=1}^M \frac{\cosh(\beta\Omega_k)}{\cosh(\beta\Omega_k) + 1} \frac{\sin^2 \theta_k \sin^2(t\Omega_k)}{\Omega_k^2}. \end{aligned} \quad (42)$$

Notice that these expressions, for the XY model, can be also obtained directly from Eq. (12). For a given λ , we can consider the time average OFS which is given by

$$\overline{\chi_{\rho}}(\lambda) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt \chi_{\rho}(\lambda, t). \quad (43)$$

Obviously $\overline{\chi_{\rho}^{(1)}}(\lambda)$ diverges, since $\chi_{\rho}^{(1)}(\lambda)$ is proportional to t^2 . The time average of the circular function contributions $\chi_{\rho}^{(2)}$ is

$$\overline{\chi_{\rho}^{(2)}}(\lambda) = 2 \sum_{k=1}^M \frac{\cosh(\beta\Omega_k)}{\cosh(\beta\Omega_k) + 1} \frac{\sin^2 \theta_k}{\Omega_k^2}. \quad (44)$$

After rescaling $\chi_{\rho} \rightarrow \chi_{\rho}/N$, $2\pi k/N \rightarrow k$ and taking the thermodynamical limit, we get

$$\begin{aligned} \chi_{\rho}^{(1)}(\lambda, t) &= \frac{2t^2}{\pi} \int_0^{\pi} dk \frac{1}{\cosh(\beta\Omega_k) + 1} \cos^2 \theta_k, \\ \chi_{\rho}^{(2)}(\lambda, t) &= \frac{2}{\pi} \int_0^{\pi} dk \frac{\cosh(\beta\Omega_k)}{\cosh(\beta\Omega_k) + 1} \frac{\sin^2 \theta_k \sin^2(t\Omega_k)}{\Omega_k^2}, \\ \overline{\chi_{\rho}^{(2)}}(\lambda) &= \frac{1}{\pi} \int_0^{\pi} dk \frac{\cosh(\beta\Omega_k)}{\cosh(\beta\Omega_k) + 1} \frac{\sin^2 \theta_k}{\Omega_k^2}. \end{aligned} \quad (45)$$




FIG. 1: First term of OFS divided by t^2 as a function of λ with different temperatures in the thermodynamical limit.

One finds that both the quantities $\chi_{\rho}^{(1)}(\lambda, t)$ and $\overline{\chi_{\rho}^{(2)}}(\lambda)$ have nontrivial behavior at the critical point




FIG. 2: Time average of the second term of OFS as a function of λ for a given $\beta = 1$ with different M .




FIG. 3: Time average of the second term of OFS as a function of λ and β for a given $M = 1000$.

$\lambda = 1$ and, therefore, can be used as indicators of criticality. At $\lambda = 1$, $\chi_{\rho}^{(1)}$ has a minimum (see Fig. 1), while $\overline{\chi_{\rho}^{(2)}}$ has a maximum, (see Fig. 3).

Notice that $\chi_{\rho}^{(1)}$ vanishes only when the inverse temperature $\beta = \infty$. When the temperature is not zero, for long enough time, χ_{ρ} will be dominated by $\chi_{\rho}^{(1)}$ and therefore will have a minimum at $\lambda = 1$. This has to be contrasted with the previous results for ground states in Ref. [5] where it was argued that the short time LE decay is enhanced at criticality (large N .) For a fixed t and λ , as temperature increases, $\chi_{\rho}^{(1)}$ grows while $\chi_{\rho}^{(2)}$ decays.

The time average of second term of OFS will diverge at the critical points $\lambda = 1$ under thermodynamical limit (see Fig. 2). It is caused by the denominator Ω_k in Eq. (45), since it will vanish for those very small k s at

the critical point $\lambda = 1$ when N goes to infinity. Thus, this divergence will be retained for all β (see Fig. 3).

In principle, one obtains $\chi_\rho^{(1)}(\lambda, t)$ and $\overline{\chi_\rho^{(2)}}(\lambda)$ from experiments in different ways. If we use $\chi_\rho^{(1)}(\lambda, t)$ to investigate the quantum critical point, the measurement of the time interval t is important. However, when $\chi_\rho^{(1)}(\lambda, t)$ vanishes, we can use $\overline{\chi_\rho^{(2)}}(\lambda)$ to investigate quantum criticality. This means we can get a $\chi_\rho^{(2)}(\lambda, t)$ for a random t again and again, then evaluate the time average value. By this scheme, we can avoid the demand of the exact measurement of the time interval t from the coupling instant to the measuring instant.

V. CONCLUSION

To summarize, we have introduced the state-dependent operator fidelity and its susceptibility χ_ρ to study environment-induced decoherence. By deriving general expressions for χ_ρ we identified two different contributions to the susceptibility. These two terms have different

physical origin and temporal behavior. For the transverse field Ising model, we obtained an exact expression for χ_ρ and showed that it has nontrivial behavior at the critical point both at zero and non-zero temperature. Moreover, from (12) it is clear that the OFS depends crucially in the level spacing distribution of H . This leads to conjecture that OFS might be an effective tool to investigate the transition to quantum chaos as well. Finally, we believe this type of analysis is directly relevant to experiments aimed at using quantum probes to detect QPTs.

VI. ACKNOWLEDGMENTS

We thank N. Toby Jacobson for a careful reading of the manuscript. The work was supported by the Program for New Century Excellent Talents in University (NCET), the NSFC with grant nos. 90503003, the State Key Program for Basic Research of China with grant nos. 2006CB921206, the Specialized Research Fund for the Doctoral Program of Higher Education with grant No.20050335087.

-
- [1] W. H. Zurek, Rev. Mod. Phys. **75**, 715-775.
 - [2] M. A. Nielsen and I. L. Chuang, *Quantum Computing and Quantum Information*, (Cambridge University Press, Cambridge, England, 2000).
 - [3] Z. K. Karkuszewski, C. Jarzynski, and W. H. Zurek, Phys. Rev. Lett. **89** 170405 (2002).
 - [4] F. M. Cucchietti, J. P. Paz, and W. H. Zurek, Phys. Rev. A **72** 052113 (2005).
 - [5] H. T. Quan, Z. Song, X. F. Liu, P. Zanardi, and C. P. Sun, Phys. Rev. Lett. **96**, 140604 (2006).
 - [6] D. Rossini et al, J. Phys. A: Math. Theor. **40**, 8033 (2007); D. Rossini et al, Phys. Rev. A **75**, 032333 (2007).
 - [7] F. M. Cucchietti, S. F. Vidal, and J. P. Paz, Phys. Rev. A **75**, 032337 (2007).
 - [8] C. Cormick, and J. P. Paz, Phys. Rev. A. **77**, 022317.
 - [9] J. Zhang, X. Peng, N. Rajendran, and D. Suter, Phys. Rev. Lett. **100**, 100501 (2008).
 - [10] R. A. Jalabert and H. M. Pastawski Phys. Rev. Lett. **86**, 2490 (2001).
 - [11] T. Gorina, T. Prosen, T. H. Seligman, and M. Žnidarič, Phys. Reports **435**, 33 (2006).
 - [12] X. Wang, Z. Sun, and Z. D. Wang, arXiv:0803.2940v2.
 - [13] P. Zanardi and N. Paunkovic, Phys. Rev. E **74**, 031123 (2006).
 - [14] H. Q. Zhou and J. P. Barjaktarevic, cond-mat/0701608.
 - [15] P. Zanardi, H. T. Quan, X. Wang, and C. P. Sun, Phys. Rev. A **75**, 032109 (2007.)
 - [16] P. Zanardi, L. C. Venuti, and P. Giorda, Phys. Rev. A **76**, 062318 (2007) .
 - [17] W. L. You, Y. W. Li, and S. J. Gu, Phys. Rev. E **76**, 022101 (2007).
 - [18] Shu Chen, Li Wang, Yajiang Hao, and Yupeng Wang Phys. Rev. A **77**, 032111 (2008).
 - [19] H. P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems*, (Oxford University Press, Oxford, 2002).
 - [20] P. Zanardi, P. Giorda, and M. Cozzini, Phys. Rev. Lett. **99**, 100603 (2007).
 - [21] L. Campos Venuti and P. Zanardi, Phys. Rev. Lett, **99**, 095701 (2007).
 - [22] E. Lieb, T. Shultz and D. Mattis, Ann. Phys. **16**, 407 (1961).
 - [23] S. Sachdev, *Quantum phase transition* (Cambridge University Press, Cambridge, England, 1999).
 - [24] P. W. Anderson, Phys. Rev. **112**, 1900 (1958).
 - [25] D. K. L. Oi, J. Aberg, Phys. Rev. Lett. **97**, 220404 (2006)